

Kunze, Tim; Hunold, Alexander; Haueisen, Jens; Jirsa, Viktor K.; Spiegler, Andreas:

Investigating the effect of electrical brain stimulation using a connectome-based brain network model

Original published in: BMC neuroscience. - London : BioMed Central. - 16 (2015), Suppl. 1, art. O13, 2 pp.
Original published: 2015-12-04
ISSN: 1471-2202
DOI: [10.1186/1471-2202-16-S1-O13](https://doi.org/10.1186/1471-2202-16-S1-O13)
[Visited: 2019-08-09]



This work is licensed under a [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

ORAL PRESENTATION

Open Access

Investigating the effect of electrical brain stimulation using a connectome-based brain network model


Tim Kunze^{1,2*}, Alexander Hunold¹, Jens Hauelsen¹, Viktor Jirsa^{3,4,5}, Andreas Spiegler^{3,4}

From 24th Annual Computational Neuroscience Meeting: CNS*2015
Prague, Czech Republic. 18-23 July 2015

Transcranial direct current stimulation (tDCS) leads to positive effects in neurological and psychiatric diseases, such as depression, pain, or stroke, which outlast the treatment itself. Although numerous influencing stimulation parameters and factors are known, the mechanisms behind tDCS remain unclear. To reveal the mechanisms tDCS started to be considered to affect networks while (de)polarizing parts of the brain. We study here the ability of tDCS as a tool to bias functional networks by affecting the connections given the brain structure.

We used structural data, that is, a human connectome to construct a large-scale brain network model of 74 cerebral areas, each described by a Jansen and Rit model. The model was designed on the basis of the neuroinformatics platform The Virtual Brain to account for reproducibility of the simulations. The tDCS-induced currents on the cerebral areas were calculated using a finite element method model. Based on the dynamical repertoire of an isolated area [1], we analyzed the brain activity, that is, the spatio-temporal dynamics in terms of rhythms and baseline potentials during rest, during tDCS, and the change between both.

We identified the network states during rest and catalogued all states for further modeling studies. During tDCS, increased functional connectivity was found among a set of scalp EEG sensors, as reported in measurements [2], as well as among cerebral cortical areas (see Figure 1). Furthermore, tDCS led to sharpened frequency spectra and increased (anode) or decreased (cathode) power in the respective areas.



* Correspondence: tim.kunze@tu-ilmenau.de

¹Institute of Biomedical Engineering and Informatics, Ilmenau University of Technology, Ilmenau, 98693, Germany

Full list of author information is available at the end of the article

This study supports the notion that noninvasive brain stimulation is able to bias brain dynamics by affecting the competitive interplay of functional subnetworks. Our work constitutes a basis for further modeling studies to test target-oriented manipulation of functional networks (e.g. through adapted electrode montages) to improve pertinent treatment conditions. Furthermore, our approach emphasizes the role of structural data such as the network topology in emerging dynamics. Dynamics cannot necessarily be predicted from the structure but we found the structure especially important at transitions of network states.

Authors' details

¹Institute of Biomedical Engineering and Informatics, Ilmenau University of Technology, Ilmenau, 98693, Germany. ²Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany. ³Aix-Marseille Université, Institut de Neurosciences des Systèmes, Marseille, France. ⁴Institut National de la Santé et de la Recherche Médicale, UMR_S 1106, 27 Bd Jean Moulin, 13385, Marseille Cedex 5, France. ⁵Centre National de la Recherche Scientifique, 3, rue Michel-Ange, 75794, Paris, France.

Published: 18 December 2015

References

1. Spiegler A, Kiebel SJ, Atay FM, Knoesche TR: **Bifurcation analysis of neural mass models: Impact of extrinsic inputs and dendritic time constants.** *Neuroimage* 2010, **52**(3):1041-1058.
2. Polania R, Nitsche MA, Paulus W: **Modulating functional connectivity patterns and topological functional organization of the human brain with transcranial direct current stimulation.** *Hum Brain Mapp* 2011, **32**(8):1236-1249.

doi:10.1186/1471-2202-16-S1-O13

Cite this article as: Kunze et al.: Investigating the effect of electrical brain stimulation using a connectome-based brain network model. *BMC Neuroscience* 2015 **16**(Suppl 1):O13.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

